



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$+ \frac{df}{dx}x + \frac{df}{dy}y + \dots + \frac{df}{dw}w, \quad (25')$$

which is of the form (25) and we have

$$\begin{aligned} r_u^2 = & \left\{ \frac{df^2}{dx^2} \left[\frac{a^2}{p} \right] + \frac{df}{dx} \cdot \frac{df}{dy} \left[\frac{a\beta}{p} \right] + \dots + \frac{df}{dx} \cdot \frac{df}{dw} \left[\frac{a\lambda}{p} \right] \right. \\ & + \frac{df}{dx} \cdot \frac{df}{dy} \left[\frac{a\beta}{p} \right] + \frac{df^2}{dy^2} \left[\frac{\beta^2}{p} \right] + \dots + \frac{df}{dy} \cdot \frac{df}{dw} \left[\frac{\beta\lambda}{p} \right] \\ & + \frac{df}{dx} \cdot \frac{df}{dw} \left[\frac{a\lambda}{p} \right] + \frac{df}{dy} \cdot \frac{df}{dw} \left[\frac{\beta\lambda}{p} \right] + \dots + \left. \frac{df^2}{dw^2} \left[\frac{\lambda^2}{p} \right] \right\} r^2. \quad (26') \end{aligned}$$

CONTINUATION OF THE NOTE AT P. 57 BY THE EDITOR.—Substituting for a its value $\frac{\omega^2 m R}{5F} + 1$, also substituting for F its value, $m \frac{v^2}{D}$, from (3), and writing h for $\omega^2 \div v^2$, we have

$$\theta = \frac{R}{r} \varphi - \frac{R}{r} \sqrt{\left(\frac{RDh}{5} \right)} \int \frac{d\varphi}{\sqrt{\left(\frac{1}{5} RDh + 1 - \cos \varphi \right)}};$$

or, substituting for $\cos \varphi$ its equivalent, $1 - 2 \sin^2 \frac{1}{2} \varphi$, dividing the quantity under the second radical by $\frac{1}{5} RDh$, placing this divisor in the denominator of the first radical and writing ψ for $\frac{1}{2} \varphi$ and c^2 for $10 \div RDh$, we get

$$\theta = \frac{R}{r} \varphi - \frac{R}{r} \int \frac{2d\psi}{\sqrt{(1 + c^2 \sin^2 \psi)}}.$$

From (3) we have

$$\begin{aligned} \theta &= \frac{R}{r} \varphi - \frac{R}{r} \omega t, \\ \therefore t &= \frac{2}{\omega} \int \frac{d\psi}{\sqrt{(1 + c^2 \sin^2 \psi)}}. \end{aligned} \quad (8)$$

Substituting for $\sin^2 \psi$ its equivalent $1 - \cos^2 \psi$, dividing the quantity under the radical by $1 + c^2$, and writing e^2 for $c^2 \div (1 + c^2)$ we have

$$t = \frac{2}{\omega \sqrt{(1 + c^2)}} \int \frac{d\psi}{\sqrt{(1 - e^2 \cos^2 \psi)}}. \quad (9)$$

The foregoing was sent to Mr. Charles H. Kummell, of the U. S. Lake Survey, with the request that he should calculate the numerical value of t ; and, in response, Mr. Kummell has put the integral in the form

$$t = \int_{\frac{1}{2}\pi - \phi}^{\frac{\pi}{2}} \frac{d\Psi}{\sqrt{(a^2 \cos^2 \Psi + b^2 \sin^2 \Psi)}} \quad (10)$$

which is more convenient for numerical calculation, and from which he finds

$$t_n = 9^h 05^m 38^s.4.$$

Hence it appears that the ball will pass a point 180° from the initial point, $2^h 54^m 21^s.6$ before the point A will pass the same point in space.